FARLOW - NOTEBOOK (?)

A
B
C
D
E
F
G
H
I
L
M
N
O
P
Q
R
S
T
U
V
X
Z
Lun
Mar
Mer

Fries. Novae
Symbolae reprint
Gen Polygoneis
Dryophila n. 66 (50)
Nova act. R. Soc.
Upsala. III 1.
In the Bibliotheque
of the Museum the
date of the volume
is 1855. But the
Fries paper is in
Swedish. Prior to which
the date 1851. The
pages are 17-136.

In Herb. De la Pylai
at Paris
My Fuciys from
Isles of Lhoals
agrees with F. Mucla
nensis in Herb. De la
Pyl. But F. Fusci
seems to be about the
same with
Zonewhat larger
and nearer F. edentatus
but not quite that
species. In Herb.
De la Pylai are
specimens labelled
F. mucronatus plus
which have marked
them F. filiformis
Gmelin in Bonnet's
handwriting. These
seem to be practically

B C D E F G IK

the same as Zacus
Mediterranean.

Largassum baccifera
in Herb Kew,
The following
localities for
other places than
W.I. and adjacent
shores.
Hobart Town
New Zealand Hombron
New Guinea "
Seychelles
Dupont
Batavia
Reynaud

Java. Perrottet.
Ceylon. Beyaard.

C D E F G

Herb. Bonet

The long Bermuda
Sargassum called
Acidendula is dut
in S. Cendigerum
a. var. Harlowii
Grunow /

Sarg. vulgare. Monahassa
Harv. = S. Cendigerum
var. Mombassiensis
Grunow.
Sarg. Eckeri. Dr Trott
Same as S. Liebmannii

Jacq. cymosum
Hohenacker 740
Areschoug Phyc. Exs.
I am correct.

Under F. cymosum
Antilles, Herb. Bory
is the note "F.
pendigerum aut
si non F. nodosa
forma brevicapsa."
Under F. pendigerum
is F. pendicymosum
are W. I. forms. This
includes F. ramifolium
Kg.

Taylor Alg. Bermuda
87, 88, attached to
stones Hamilton

is Sargassum Cini
folium var. Bermudense
Grun. No. 85 on
rocks at Low tide
South Beach is
the same forma
contracta Grun.

There is a S. fibuliden-
data var. subcomosum
Grun from Brinjolvd
in Heib. Bornet.

The Sargassum
cymosum Zat
Lauderdale Florida
a. H. Curtiss April 1897
Not in S. vulgare.

C
D
E
F
G
in

Herbarium 434
S. tracht. phyllum
N.D. as S. pulgare
Poiosis 2 cm.

D
E
F
G







D
E
F
G

Herb. Barnet Jul 7.

Lang. Lendigerum
var. pseudocymosum
Gren. fama furcata
Gren. Bor. Martinique
marked "ad S. Vulgarem
Leus - minus ne accedit.

Lang. Turneri Crn.
(Cum a cultus Hz.)
Grenoble Combe is
S. Matycarpum Mart.
Dec. Gren. Is there
a specimen from
Vickers of this in my
herbarium.

Then is the following
note in Herb. Barnet,
"Lucelle Chaine,
S. Matycarpum,

spinulosum, Esseni
Lonchitis (Retzium)?
(Retzium)?
L. Boottii
spinulosum Lise
L. Leebmannii?

L. vulgare

L. diversifolium Ky.
Neuz. Schramm.
Just. Ky. Ennom. in
L. vulgare v. diversifolium
L. megalophyllum
mont. rar. f.
vulgare. forma in
alpinis.
L. trachyphyllum
Hohenacker 43 d
L. vulgare. foliorum

L. Bahyense =
L. nutans foliosissimum
also L. cheiri folium
L. Cestocarpum
forma elegans. Tortola
L. micranthum
Parallela Cuba
= foliosissimum

L. Hystrix in Herb.
Barnet. Cult. but
very doubtful
Specimens

L. Hystrix in Herb.
Pers. one specimen
from Bahama given
by Descaisne 1849.
marked L. Thom-
asyi n. sp. This
specimen not at
all characteristic

Herb. Paris

I. bacciferum
angustum. Collins
Berg Ged

I. cymosum
includes I. Esperi
Martindyce

The specimens of
I. cymosum, espec-
ially I. stenophyllum
are almost all
from Brazil
I. lendigerum in Herb
Paris from Red Sea
Cape Verds, Brazil
but now from Kew

VI